

AN INDEX THEOREM FOR GAUGE-INVARIANT FAMILIES: THE CASE OF SOLVABLE GROUPS

VICTOR NISTOR

Version: 1.2; Revised: August 7, 2001; Run: February 1, 2008

ABSTRACT. We define the gauge-equivariant index of a family of elliptic operators invariant with respect to the free action of a family $\mathcal{G} \rightarrow B$ of Lie groups (these families are called “gauge-invariant families” in what follows). If the fibers of $\mathcal{G} \rightarrow B$ are simply-connected and solvable, we compute the Chern character of the gauge-equivariant index, the result being given by an Atiyah-Singer type formula that incorporates also topological information about the bundle $\mathcal{G} \rightarrow B$. The algebras of invariant pseudodifferential operators that we study, $\psi_{\text{inv}}^\infty(Y)$ and $\Psi_{\text{inv}}^\infty(Y)$, are generalizations of “parameter dependent” algebras of pseudodifferential operators (with parameter in $\mathbb{R}^d$), so our results provide also an index theorem for elliptic, parameter dependent pseudodifferential operators. We apply these results to study Fredholm boundary conditions on a simplex.

CONTENTS

Introduction	1
1. Invariant families of (pseudo)differential operators	4
2. The case when $\mathcal{G}$ is a vector bundle	6
3. The gauge-equivariant index: Definition	8
4. The gauge-equivariant index: A formula	11
5. Index theory on a simplex	20
References	22

INTRODUCTION

Families of operators invariant with respect to the action of a family of Lie groups (referred to as “gauge-invariant families,” in what follows) have been considered in several papers in the literature. They appear, for example, in the work of Bismut on analytic torsion [9], in the work of Connes on abstract index theory on groupoids [14], and in the work of Mazzeo and Melrose on analysis on non-compact manifolds [23, 24, 25]. Gauge-invariant operators are also a basic example of the operators on groupoids introduced in [37].

Gauge-invariant families are likely to form the main building block in the analysis of operators on a certain class of non-compact manifolds (manifolds with a Lie

Partially supported by the NSF Young Investigator Award DMS-9457859, a Sloan fellowship, and NSF Grant DMS-9971951. <http://www.math.psu.edu/nistor/>.

structure at infinity [2, 19, 21, 25, 27]). This class of non-compact manifolds is important for the analysis on locally symmetric spaces [31, 32].

Our main motivation to study gauge-invariant families comes from analysis and spectral theory on non-compact manifolds. For example, the Fredholmness of an elliptic differential operator on a manifold with a uniform structure at infinity is controlled by (possibly several) gauge-invariant families of elliptic differential operators, see [18, 20, 21]. This is a situation that arises when one studies the Dirac operator on an S^1 -manifold M , if we desingularize the action of S^1 by replacing the original metric g with $\phi^{-2}g$, where ϕ is the length of the infinitesimal generator X of the S^1 -action. In this way, X becomes of length one in the new metric. The main result of [35] states that the kernel of the new Dirac operator on the open manifold $M \setminus M^{S^1}$ is naturally isomorphic to the kernel of the original Dirac operator. It turns out that the Fredholm property of the resulting Dirac operator (obtained by the above procedure on the non-compact manifold $M \setminus M^{S^1}$) is controlled by the invertibility of a family of operators invariant with respect to the action of a bundle of non-abelian, solvable Lie groups.

In this paper, we study the gauge-equivariant index of families of elliptic operators invariant with respect to the free action a bundle of *simply-connected, solvable* Lie groups $\mathcal{G}$. (By “simply-connected” we shall mean, as usual, “connected with trivial fundamental group.”) This is the setting derived from the example of the S^1 -Dirac operator considered above and is enough in many applications. The case of a bundle of compact Lie groups will be treated in a forthcoming paper with E. Troitsky [36]. We are interested in the gauge-equivariant index of invariant families because the non-vanishing of the gauge-equivariant index is a first obstruction to the invertibility of that family. One of the main results of this paper is a formula for the Chern character of the gauge-equivariant index that is similar to the Atiyah-Singer index formula for families, but also includes information on the *topology* of the bundle of Lie groups that is considered, which is a qualitatively new feature of our setting.

We define the gauge-equivariant index of a family of invariant, elliptic operators using K -theory. Our definition works for arbitrary family of Lie groups, provided that they act freely. (Both a *bundle of Lie groups* and a *family of Lie groups* are fiber bundles with homeomorphic fibers. For a bundle of Lie groups the fibers are required to be isomorphic as Lie groups. This is not required, however, for families of Lie groups.)

In [10], Bismut and Cheeger have generalized the Atiyah-Patodi-Singer index theorem [6] to families of Dirac operators on manifolds with boundary (see also [29]). Their results apply to operators whose “indicial parts” are invertible. These indicial parts are actually families of Dirac operators invariant with respect to a one-parameter group, so they fit into the framework of this paper (with $\mathcal{G} = B \times \mathbb{R}$). Thus, the results of this paper are relevant also to the problem of studying families of operators on fibrations by manifolds with boundary or fibrations by more general singular spaces. (See also [44].)

We now describe the contents of each section of this paper. In Section 1, we discuss the action of a bundle of Lie groups $\mathcal{G} \rightarrow B$ on a fiber bundle $Y \rightarrow B$ and we introduce the algebras $\psi_{\text{inv}}^\infty(Y)$ of families of invariant pseudodifferential operators on the fibers of $Y \rightarrow B$ whose convolution kernels are compactly supported. The classical definition of symbols applies to algebras $\psi_{\text{inv}}^\infty(Y)$ also. In Section 2 and

thereafter we shall assume that $\mathcal{G}$ acts freely on Y . Section 2 is devoted to the case when $\mathcal{G}$ consists of simply-connected, abelian Lie groups (which is the same thing as saying that $\mathcal{G} \rightarrow B$ is a vector bundle, with the Lie group structure on each fiber given simply by the addition of vectors). We prove then that the group of gauge transformations of $\mathcal{G}$ acts on $\psi_{\text{inv}}^\infty(Y)$, and we use this to enlarge the algebras $\psi_{\text{inv}}^\infty(Y)$ by including families whose convolution kernels are Schwartz functions. The resulting algebras will be denoted by $\Psi_{\text{inv}}^\infty(Y)$. Let $C_r^*(\mathcal{G})$ be the closure of $\psi_{\text{inv}}^\infty(Y)$ in the topology defined by its action on $L^2(Y)$. In Section 3, we consider arbitrary families of Lie groups $\mathcal{G} \rightarrow B$, but continue to assume that they act freely on Y . We define the gauge-equivariant index

$$\text{ind}_{\mathcal{G}}(A) \in K_0(C_r^*(\mathcal{G}))$$

of a family of elliptic, invariant pseudodifferential operators $A \in \psi_{\text{inv}}^\infty(Y)$ (or $A \in \Psi_{\text{inv}}^\infty(Y)$, if $\mathcal{G}$ is a vector bundle). We then prove that the gauge-equivariant index of A is the obstruction to finding a regularizing family R such that $A + R$, acting between suitable Sobolev spaces, is invertible in each fiber, provided that $\dim Y > \dim \mathcal{G}$. This generalizes the usual property of the Fredholm index of an elliptic pseudodifferential operator. In Section 4, we consider bundles of simply-connected, solvable Lie groups $\mathcal{G} \rightarrow B$, and we prove that

$$(1) \quad K_*(C_r^*(\mathcal{G})) \simeq K^*(\mathfrak{g}),$$

provided that $\mathcal{G}$ consists of simply-connected solvable Lie groups, where $\mathfrak{g} := \cup \text{Lie } \mathcal{G}_b \rightarrow B$ is the vector bundle of Lie algebras defined by $\mathcal{G}$. Then we obtain a formula for the Chern character of the gauge-equivariant index of an elliptic operator $A \in \psi_{\text{inv}}^\infty(Y)$. (For simplicity, we called A “an operator,” although it is really a family of operators. We shall do this repeatedly.) The proof of this result requires us to construct a non-trivial deformation of the indicial family to a family to which the Atiyah-Singer index theorem for families can be applied. We also use here the notion of “degree of an elliptic family.”

Operators invariant with respect to $\mathbb{R}^q$, a particular case of our operators when the base is reduced to a point, appear in the formulation of elliptic (or Fredholm) boundary conditions for pseudodifferential operators on manifolds with corners. The gauge-equivariant index can be used to study this problem, which is relevant to the question of extending the Atiyah-Patodi-Singer boundary conditions to manifolds with corners. This is discussed in the last section.

The algebras of invariant pseudodifferential operators that we study, $\psi_{\text{inv}}^\infty(Y)$ and $\Psi_{\text{inv}}^\infty(Y)$, are generalizations of “parameter dependent” algebras of pseudodifferential operators considered by Agmon [1], Grubb and Seeley [16], Lesch and Pflaum [22], Melrose [27], Shubin [41], and others. Our index theorem, Theorem 5, thus gives a solution to the problem of determining the index of elliptic, parameter dependent families of pseudodifferential operators, parameterized by $\lambda \in \mathbb{R}^q$. We note that the concept of index of such a family requires a proper definition, and that the Fredholm index, “dimension of the kernel” - “dimension of the cokernel,” is not adequate for $q \geq 1$. Actually, our definition of the gauge-equivariant index is somewhat closer to the definition of the L^2 -index for covering spaces given in [4] and [40] than to the definition of the Fredholm index. However, an essential difference between their definition and ours is that no trace is involved in our definition.

There exist several potential extensions of our results, although none of them seems to be straightforward. These extensions will presumably involve more general

conditions on $\mathcal{G}$ and its action on Y . Probably the most general conditions under which one can reasonably expect to obtain definite results are those when the fibers of $\mathcal{G}$ are connected and the action is proper. We need the fibers of $\mathcal{G}$ to be connected in order to be able to use general results on the structure of connected Lie groups. The case when the fibers of $\mathcal{G} \rightarrow B$ are compact Lie groups will be discussed in [36], where the relevance of these construction to the study of Ramond-Ramond charges and of the Dixmier-Douady-Donovan-Karoubi invariant.

We plan to apply the results of this paper to some problems in M -theory [39], to the study of adiabatic limits of eta invariants [11, 45], and to analysis on locally-symmetric spaces [31, 32].

I would like to thank Fourier Institute and University of Grenoble, where part of this project was completed, for their hospitality. Also, I am grateful to Sergiu Moroianu and Evgenij Troitsky for useful discussions. A preliminary version of this paper was circulated as a Fourier Institute preprint in 1999.

All pseudodifferential operators considered in this paper are “classical,” that is, one-step polyhomogeneous.

1. INVARIANT FAMILIES OF (PSEUDO)DIFFERENTIAL OPERATORS

We now describe the settings in which we shall work, that of a family of Lie groups $p : \mathcal{G} \rightarrow B$ acting on a fiber bundle $\pi : Y \rightarrow B$. Then we introduce the algebra $\psi_{\text{inv}}^\infty(Y)$ consisting of families of pseudodifferential operators along the fibers of Y that are invariant under the action of $\mathcal{G}$ and whose convolution kernels are compactly supported. These algebras are particular cases of the algebras introduced in [37]. We can restrict in our discussion to a connected component of B , so for simplicity, we shall assume that B is connected. The gauge-equivariant index of the elliptic operators $A \in \psi_{\text{inv}}^\infty(Y)$ will form our main object of study.

Throughout this paper, $p : \mathcal{G} \rightarrow B$ will denote a *family of Lie groups* on a smooth manifold B . By this we mean that $p : \mathcal{G} \rightarrow B$ is a smooth fiber bundle, that each $\mathcal{G}_b := p^{-1}(b)$ is a Lie group, and that the multiplication and inverse depend differentiably on b . Hence the map

$$(2) \quad \mathcal{G} \times_B \mathcal{G} := \{(g', g) \in \mathcal{G} \times \mathcal{G}, d(g') = d(g)\} \ni (g', g) \longrightarrow g'g^{-1} \in \mathcal{G}$$

is differentiable. This implies, by standard arguments, that the map sending a point $b \in \mathcal{G}_b := p^{-1}(b)$ to e_b , the identity element of $\mathcal{G}_b$, is a diffeomorphism onto a smooth submanifold of $\mathcal{G}$. It also implies that the map $\mathcal{G} \ni g \rightarrow g^{-1} \in \mathcal{G}$ is differentiable. Note that we do not assume the groups $\mathcal{G}_b$ to be isomorphic, although this is true in most applications. Let $\mathfrak{g} := \cup \text{Lie } \mathcal{G}_b \rightarrow B$ be the bundle of Lie algebras of associated to $\mathcal{G}$, that is, the bundle whose fiber above b , $\mathfrak{g}_b$, is $\text{Lie } \mathcal{G}_b$.

Let us remark, although we shall not use this here, that $\mathcal{G}$ is a particular case of a differentiable groupoid. The Lie algebroid associated to this groupoid is the bundle $\mathfrak{g} \rightarrow B$ defined above. (See [37] or [21] for the necessary definitions.)

We also assume that $\mathcal{G}$ acts smoothly on a fiber bundle $\pi : Y \rightarrow B$. This means that there are given actions $\mathcal{G}_b \times Y_b \rightarrow Y_b := \pi^{-1}(b)$ of $\mathcal{G}_b$ on Y_b , for each b , such that the induced map,

$$\mathcal{G} \times_B Y := \{(g, y) \in \mathcal{G} \times Y, d(g) = \pi(y)\} \ni (g, y) \longrightarrow gy \in Y,$$

is differentiable. The action of $\mathcal{G}$ on Y is called *free* [respectively, *proper*], if the action of $\mathcal{G}_b$ on Y_b is free [respectively, proper] for each b . (In this section we make

no assumptions on the action of $\mathcal{G}$ on Y , other than that it is smooth, but beginning with the next section we shall assume that this action is free.)

On Y , we consider smooth families $A = (A_b)$, $b \in B$, of classical pseudodifferential operators acting on the fibers of $Y \rightarrow B$. Unless mentioned otherwise, we assume that these operators act on half densities along each fiber. The family $(A_b)_{b \in B}$ will be called $\mathcal{G}$ -invariant if each A_b is invariant with respect to the action of the group $\mathcal{G}_b$. A *gauge-invariant family* is a $\mathcal{G}$ -invariant family, for some family of Lie groups $\mathcal{G}$. The algebra that we are interested in consists of $\mathcal{G}$ -invariant operators satisfying also the following *support condition*. To state this support condition, notice first that a family $A = (A_b)$ defines a continuous map $\mathcal{C}_c^\infty(Y) \rightarrow \mathcal{C}^\infty(Y)$, and, as such, it has a distribution (or Schwartz) kernel, which is a distribution K_A on $Y \times_B Y \subset Y \times Y$. (We ignore the vector bundles in which this distribution takes its values.) Because the family $A = (A_b)$ is invariant, the distribution K_A is also invariant with respect to the action of $\mathcal{G}$. Consequently, the distribution K_A is the pull back of a distribution k_A defined on $(Y \times_B Y)/\mathcal{G}$. We will require that k_A have compact support. We shall sometimes call k_A the *convolution kernel* of A . This condition on the support of k_A ensures that each A_b is a properly supported pseudodifferential operator, and hence it maps compactly supported functions (or sections of a vector bundle, if we consider operators acting on sections of a smooth vector bundle) to compactly supported functions (or sections). This support condition is automatically satisfied if $Y/\mathcal{G}$ is compact and each A_b is a differential operator. The space of smooth, invariant families A of order m pseudodifferential operators acting on the fibers of $Y \rightarrow B$ such that k_A has compact support will be denoted by $\psi_{\text{inv}}^m(Y)$. Then

$$\psi_{\text{inv}}^\infty(Y) := \cup_{m \in \mathbb{Z}} \psi_{\text{inv}}^m(Y)$$

is an algebra, by classical results [17, 42, 43]. Note also that $\psi_{\text{inv}}^m(Y)$ makes sense also for m not an integer.

We now discuss the principal symbols of an operator $A \in \psi_{\text{inv}}^m(Y)$. Let

$$T_{\text{vert}}Y := \ker(TY \rightarrow TB)$$

be the bundle of *vertical* tangent vectors to Y , and let T_{vert}^*Y be its dual. We fix compatible metrics on $T_{\text{vert}}Y$ and T_{vert}^*Y , and define S_{vert}^*Y , the *cosphere bundle of the vertical tangent bundle* to Y , to be the set of vectors of length one of T_{vert}^*Y . Also, let

$$\sigma_m : \Psi^m(Y_b) \rightarrow \mathcal{C}^\infty(S_{\text{vert}}^* \cap T^*Y_b)$$

be the usual principal symbol map, defined on the space of pseudodifferential operators of order m on Y_b . The definition of σ_m depends on the choice of a trivialization of the bundle of homogeneous functions of order m on T_{vert}^*Y , regarded as a bundle over S_{vert}^*Y . The principal symbols $\sigma_m(A_b)$ of an element (or family) $A = (A_b) \in \psi_{\text{inv}}^m(Y)$ then gives rise to a smooth function on $\mathcal{C}^\infty(S_{\text{vert}}^*Y)$, which is invariant with respect to $\mathcal{G}$, and hence descends to a smooth function on S_{vert}^*Y . Let $\mathcal{C}^\infty(S_{\text{vert}}^*Y)^\mathcal{G}$ be the space of smooth functions on S_{vert}^*Y that are invariant with respect to the canonical action of $\mathcal{G}$. The resulting function,

$$(3) \quad \sigma_m(A) \in \mathcal{C}^\infty(S_{\text{vert}}^*Y)^\mathcal{G},$$

will be referred to as the *principal symbol* of an element (or operator) in $\psi_{\text{inv}}^m(Y)$.

In the particular case $Y = \mathcal{G}$, $\psi_{\text{inv}}^\infty(\mathcal{G})$ identifies with families of convolution operators on the fibers $\mathcal{G}_b$ with kernels contained in a compact subset of $\mathcal{G}$, smooth

outside the identity, and only with conormal singularities at the identity. In particular, $\psi_{\text{inv}}^{-\infty}(\mathcal{G}) = \mathcal{C}_c^\infty(\mathcal{G})$, with the fiberwise convolution product.

Suppose now that the quotient $Y/\mathcal{G}$ is compact, which implies that $(S_{\text{vert}}^*Y)/\mathcal{G}$ is also compact. As is customary, an operator $A \in \psi_{\text{inv}}^m(Y)$ is called *elliptic* if, and only if, its principal symbol is everywhere invertible. The same definition applies to $A = [A_{ij}] \in M_N(\psi_{\text{inv}}^m(Y))$. Namely, the operator A , regarded as acting on sections of the trivial vector bundle $\mathbb{C}^N$, is elliptic if, and only if, its principal symbol

$$\sigma_m(A) := [\sigma_m(A_{ij})] \in M_n(\mathcal{C}^\infty(S_{\text{vert}}^*Y)^{\mathcal{G}})$$

is invertible.

Assume that there is given a $\mathcal{G}$ -invariant metric on $T_{\text{vert}}Y$, the bundle of vertical tangent vectors, and a $\mathcal{G}$ -equivariant bundle W of modules over the Clifford algebras of $T_{\text{vert}}Y$. Then a typical example of a family $D = (D_b) \in \psi_{\text{inv}}^\infty(Y)$ is that of the family of Dirac operators D_b acting on the fibers Y_b of $Y \rightarrow B$. (Each D_b acts on sections of $W|_{Y_b}$, the restriction of the given Clifford module W to that fiber.)

The considerations of this section extend immediately to operators acting between sections of two $\mathcal{G}$ -equivariant vector bundles. If E_0 and E_1 are $\mathcal{G}$ -equivariant vector bundles, we denote by $\psi_{\text{inv}}^m(Y; E_0, E_1)$ the space of $\mathcal{G}$ -invariant families of order m pseudodifferential operators acting on sections of E_0 , with values sections of E_1 , whose convolution kernels have compact support.

2. THE CASE WHEN $\mathcal{G}$ IS A VECTOR BUNDLE

Beginning with this section and throughout the rest of this paper, we shall assume that the action of $\mathcal{G} \rightarrow B$ on $Y \rightarrow B$ is free (that is, that the action of $\mathcal{G}_b$ on Y_b is free, for any $b \in B$). In this section we shall also assume that the fibers of $\mathcal{G} \rightarrow B$ are simply-connected, abelian Lie groups. This assumption is equivalent to the assumption that $\mathcal{G} \rightarrow B$ is a vector bundle with the induced bundle of Lie algebras structure.

For the discussion of the gauge-equivariant index of an operator $A \in \psi_{\text{inv}}^m(Y)$ in Section 3, we need to introduce the algebra $\Psi_{\text{inv}}^\infty(Y)$, which is a variant of the algebra $\psi_{\text{inv}}^\infty(Y)$ which it contains. The reason for considering the slightly larger algebra $\Psi_{\text{inv}}^\infty(Y)$ is that it will have the following property, if $A = (A_b) \in \Psi_{\text{inv}}^\infty(Y)$ and each A_b is invertible on $L^2(Y_b)$ (in the sense of unbounded operators), then $(A_b^{-1}) \in \Psi_{\text{inv}}^\infty(Y)$. (This property of the algebra $\Psi_{\text{inv}}^\infty(Y)$ is usually referred to in the literature as “spectral invariance.”)

Take $Y = B \times Z \times \mathbb{R}^q$, with Z a compact manifold and $\mathcal{G} = B \times \mathbb{R}^q$, $\pi : Y \rightarrow B$ and $p : \mathcal{G} \rightarrow B$ being the projections onto the first components of each product. The action of $\mathcal{G}$ on Y is given by translation on the last component of Y . Then the $\mathcal{G}$ -invariance condition becomes simply $\mathbb{R}^q$ invariance with respect to the resulting $\mathbb{R}^q$ action. If Y and $\mathcal{G}$ are as described here, then we call Y a *flat $\mathcal{G}$ -space*. The construction of the algebra $\Psi_{\text{inv}}^\infty(Y)$ is local, so we may assume that Y is a flat $\mathcal{G}$ -space.

The residual ideal of the algebra $\psi_{\text{inv}}^\infty(Y)$ is $\psi_{\text{inv}}^{-\infty}(Y)$ and consists of operators that are regularizing along each fiber. More precisely, it consists of those families of smoothing operators on $Y = B \times Z \times \mathbb{R}^q$ that are translation-invariant under the action of $\mathbb{R}^q$ and have *compactly supported* convolution kernels. Thus

$$\psi_{\text{inv}}^{-\infty}(Y) \cong \mathcal{C}_c^\infty(B \times \mathbb{R}^q; \Psi^{-\infty}(Z)) \subset \mathcal{S}(B \times \mathbb{R}^q; \Psi^{-\infty}(Z)) \simeq \mathcal{S}(B \times Z \times Z \times \mathbb{R}^q).$$

(Here $\mathcal{S}$ is the generic notation for the space of Schwartz functions.) The second isomorphism above is obtained from the isomorphism

$$\Psi^{-\infty}(Z) \simeq \mathcal{C}^\infty(Z \times Z),$$

defined by the choice of a nowhere vanishing density on Z . If we also endow $\Psi^{-\infty}(Z)$ with the locally convex topology induced by this isomorphism, then it becomes a nuclear, locally convex space. We now enlarge the algebra $\psi_{\text{inv}}^\infty(Y)$ to include all $\mathcal{G}$ -invariant, regularizing operators whose kernels are in $\mathcal{S}(B \times Z \times Z \times \mathbb{R}^q)$:

$$(4) \quad \Psi_{\text{inv}}^m(Y) = \psi_{\text{inv}}^m(Y) + \mathcal{S}(B \times Z \times Z \times \mathbb{R}^q).$$

Explicitly, the action of $T \in \mathcal{S}(B \times Z \times Z \times \mathbb{R}^q)$ on a smooth function $f \in \mathcal{C}_c^\infty(B \times Z \times \mathbb{R}^q)$ is given by

$$Tf(b, y_1, t) = \int_{Z \times \mathbb{R}^q} T(b, y_1, y_0, t - s) f(b, y_0, s) dy_0 ds.$$

For B reduced to a point, the algebras $\Psi_{\text{inv}}^\infty(Y)$ were considered before (see [27] and the references therein) as the range space of the indicial map for “cusp” pseudodifferential operators on manifolds with corners

Still assuming that we are in the case of a flat $\mathcal{G}$ -space, we notice that the Fourier transformation $\mathcal{F}_2$ in the translation-invariant directions gives a dual identification

$$(5) \quad \Psi_{\text{inv}}^{-\infty}(Y) \ni T \rightarrow \hat{T} := \mathcal{F}_2 T \mathcal{F}_2^{-1} \in \mathcal{S}(B \times Z \times Z \times \mathbb{R}^q) = \mathcal{S}(B \times \mathbb{R}^q; \Psi^{-\infty}(Z))$$

with the space of Schwartz functions with values in the smoothing ideal of Z . The point of this identification is that the convolution product is transformed into the pointwise product. The Schwartz topology from (5) then gives $\Psi_{\text{inv}}^{-\infty}(Y)$ the structure of a nuclear locally convex topological algebra. Following the same recipe, the Fourier transform also gives rise to an *indicial family*

$$(6) \quad \Phi : \Psi_{\text{inv}}^\infty(Y) \ni T \rightarrow \hat{T} := \mathcal{F}_2 T \mathcal{F}_2^{-1} \in \mathcal{C}^\infty(B \times \mathbb{R}^q; \Psi^\infty(Z)).$$

We denote $\hat{T}(\tau) = \Phi(T)(\tau)$.

The map Φ is not an isomorphism since $\hat{A}(\tau)$ has joint symbolic properties in the variables of $\mathbb{R}^q$ and the fiber variables of T^*Z . Actually, the principal symbols of the operators $\hat{A}(\tau)$, for τ in a fixed fiber of $T_{\text{vert}}^*Y \rightarrow B$, is independent of τ .

Lemma 1. *Assume $Y = B \times Z \times \mathbb{R}^q$ is a flat $\mathcal{G}$ -space. Then the action of the group $GL_q(\mathbb{R})$ on the last factor of $Y = B \times Z \times \mathbb{R}^q$ extends to an action by automorphisms of $\mathcal{C}^\infty(B, GL_q(\mathbb{R}))$ on $\psi_{\text{inv}}^\infty(Y)$ and on $\Psi_{\text{inv}}^\infty(Y)$.*

Proof. The vector representation of $GL_q(\mathbb{R})$ on the second component of $Z \times \mathbb{R}^q$ defines an action of $GL_q(\mathbb{R})$ on $\Psi^\infty(Z \times \mathbb{R}^q)$ that preserves the class of properly supported operators and the products of such operators. It also normalizes the group $\mathbb{R}^q$ of translations, and hence it maps $\mathbb{R}^q$ -invariant operators to $\mathbb{R}^q$ -invariant operators. This property extends right away to the action of $\mathcal{C}^\infty(B, GL_q(\mathbb{R}))$ on families of operators on $B \times Z \times \mathbb{R}^q$, and hence $\mathcal{C}^\infty(B, GL_q(\mathbb{R}))$ maps $\psi_{\text{inv}}^m(Y)$ isomorphically to itself. From the isomorphism (5), we see that $\mathcal{C}^\infty(B, GL_q(\mathbb{R}))$ also maps $\Psi_{\text{inv}}^{-\infty}(Y)$ to itself. This gives an action by automorphisms of $\mathcal{C}^\infty(B, GL_q(\mathbb{R}))$ on $\Psi_{\text{inv}}^\infty(Y)$, which is the sum of $\Psi_{\text{inv}}^{-\infty}(Y)$ and $\psi_{\text{inv}}^\infty(Y)$. $\square$

By choosing a lift of $Y/\mathcal{G} \rightarrow Y$, which is possible because the fibers are contractable ($\mathcal{G}$ is a vector bundle), we obtain that, locally, the bundle Y is isomorphic

to a flat $\mathcal{G}$ space. Then the above lemma allows us to extend the previous definitions, including those of the algebras $\Psi_{\text{inv}}^\infty(Y)$ and of the indicial family from the flat case to the case $\mathcal{G}$ a vector bundle. Recall the bundle $\mathfrak{g} := \cup \text{Lie } \mathcal{G}_b \rightarrow B$. The indicial family $\hat{A}$ of an operator $A \in \Psi_{\text{inv}}^\infty(Y)$, will then be a family of pseudodifferential operators acting on the fibers of $Y \times_B \mathfrak{g}^* \rightarrow \mathfrak{g}^*$ (here $\mathfrak{g}^*$ is the dual of the vector bundle $\mathfrak{g}$):

$$(7) \quad \hat{A}(\tau) \in \Psi^*(Y_b/\mathcal{G}_b), \quad \text{if } \tau \in \mathfrak{g}_b^* := (\text{Lie } \mathcal{G}_b)^*.$$

The action of $GL_q(\mathbb{R})$ in the above lemma will have to be replaced with the group of gauge-transformations of $\mathfrak{g}$.

We can look at a general vector bundle $\mathcal{G}$ from an other, related point of view. Let $Z' := Y/\mathcal{G}$, which is again a fiber bundle $Z' \rightarrow B$. The algebra $\Psi^\infty(Z'|B)$ of smooth families of pseudodifferential operators along the fibers of $Z' \rightarrow B$ can be regarded as the algebra of sections of a vector bundle on Z' (with infinite dimensional fibers). We can lift the vector bundle associated $\Psi^\infty(Z'|B)$ to $\mathfrak{g}^*$ and then $\hat{A}$ is a section of this lifted vector bundle over $\mathfrak{g}^*$. We shall write this as follows:

$$(8) \quad \hat{A} \in \Gamma(\mathfrak{g}^*, \Psi^\infty(Z'|B)).$$

3. THE GAUGE-EQUIVARIANT INDEX: DEFINITION

In this section, $\mathcal{G} \rightarrow B$ will an arbitrary family of Lie groups (we shall make no assumptions on the fibers $\mathcal{G}_b$), but we shall continue to assume that the action of $\mathcal{G}$ on Y is free

We now define study $\mathcal{G}$ -invariants of elliptic operators in $M_N(\psi_{\text{inv}}^\infty(Y))$, the main invariant being the gauge-equivariant index of such an invariant, elliptic family. If $\dim Y > \dim \mathcal{G}$, we then show that the gauge-equivariant index gives the obstruction for family $A = (A_b) \in M_N(\psi_{\text{inv}}^m(Y))$ to have a perturbation by a family $R = (R_b) \in M_N(\psi_{\text{inv}}^{-\infty}(Y))$, such that $A + R = (A_b + R_b)$ be invertible, for all $b \in B$, between suitable Sobolev spaces, see Theorem 1. For families of simply-connected, abelian Lie groups $\mathcal{G}$, we give an interpretation of the gauge-equivariant index of an elliptic operator in terms of its indicial family. This requires the notion of “degree,” due to Melrose. We shall use this in the next section to prove an Atiyah-Singer index type formula for the Chern character of the index of a family of $\mathcal{G}$ -invariant, elliptic operators. If $\mathcal{G}$ is a vector bundle, then we can consider the algebra $\Psi_{\text{inv}}^\infty(Y)$ instead of $\psi_{\text{inv}}^\infty(Y)$.

We now proceed to define the gauge-equivariant index of an elliptic family $A \in \psi_{\text{inv}}^m(Y)$. This will be done using the K -theory of Banach algebras. Let $C_r^*(Y, \mathcal{G})$ be the closure of $\psi_{\text{inv}}^{-\infty}(Y)$ with respect to the norm

$$\|A\| = \sup_{b \in B} \|A_b\|$$

each operator A_b acting on the Hilbert space of square integrable densities on the fiber Y_b . If $Y = \mathcal{G}$, then we also write $C_r^*(\mathcal{G}, \mathcal{G}) = C_r^*(\mathcal{G})$. (We remark, for those familiar with the concepts, that $C_r^*(\mathcal{G})$ is the reduced C^* -algebra associated to the groupoid $\mathcal{G}$. Also, the full and the reduced C^* -algebras, $C^*(\mathcal{G})$ and $C_r^*(\mathcal{G})$ coincide, if the fibers $\mathcal{G}_b$ are solvable groups.) For each locally compact space X , we denote by $C_0(X)$ the space of continuous functions vanishing at infinity on X . Then, if $\mathcal{G}$ is a vector bundle, we have $C_r^*(\mathcal{G}) \simeq C_0(\mathcal{G}^*)$.

We shall use below $\hat{\otimes}$, the minimal tensor product of C^* -algebras. This minimal tensor product is defined to be (isomorphic to) the completion of the image of

$\pi_1 \otimes \pi_2$, the tensor product of two injective representations π_1 and π_2 . For the cases we are interested in, the minimal and the maximal tensor product coincide [38].

Lemma 2. *Assume as above that $\mathcal{G}$ acts freely on Y and that $\dim Y > \dim \mathcal{G}$. Also, let $\mathcal{K} = \mathcal{K}(Y_b/\mathcal{G}_b)$ denote the algebra of compact operators on one of the fibers $Y_b/\mathcal{G}_b$, for some fixed but arbitrary $b \in B$. Then*

$$C_r^*(Y, \mathcal{G}) \simeq C_r^*(\mathcal{G}) \hat{\otimes} \mathcal{K}.$$

Consequently, $K_i(C_r^*(Y, \mathcal{G})) \simeq K_i(C_r^*(\mathcal{G}))$.

Proof. Let $\mathfrak{A}$ be the space of sections of the bundle of algebras $\mathcal{K}(Y_b/\mathcal{G}_b)$. If Y is a flat $\mathcal{G}$ -space, then the isomorphism $C_r^*(Y, \mathcal{G}) \simeq C_r^*(\mathcal{G}) \hat{\otimes} \mathcal{K}$ follows, for example, from the results of [18]. In general, this local isomorphism gives $C_r^*(Y, \mathcal{G}) \simeq C_r^*(\mathcal{G}) \hat{\otimes}_{C_0(B)} \mathfrak{A}$.

Our assumptions imply that $\mathcal{K}$ is infinite dimensional, and hence its group of automorphisms is contractable, see [15]. Consequently, there is no obstruction to trivialize the bundle of algebras $\mathcal{K}(Y_b/\mathcal{G}_b)$, and hence $\mathfrak{A} \simeq C_0(B) \hat{\otimes} \mathcal{K}$. We obtain

$$C_r^*(Y, \mathcal{G}) \simeq C_r^*(\mathcal{G}) \hat{\otimes}_{C_0(B)} \mathfrak{A} \simeq C_r^*(\mathcal{G}) \hat{\otimes}_{C_0(B)} [C_0(B) \hat{\otimes} \mathcal{K}] \simeq C_r^*(\mathcal{G}) \hat{\otimes} \mathcal{K}.$$

The last part of lemma follows from the above isomorphisms and from the natural isomorphism $K_i(A \hat{\otimes} \mathcal{K}) \simeq K_i(A)$, valid for any C^* -algebra A . $\square$

We proceed now to define the gauge-equivariant index of an *elliptic, $\mathcal{G}$ -invariant family* of operators

$$A = (A_b) \in M_N(\psi_{\text{inv}}^m(Y)) = \psi_{\text{inv}}^m(Y; \mathbb{C}^N)$$

when $Y/\mathcal{G}$ is compact. More general operators can be handled similarly, and the necessary changes will be discussed below. We observe first that there exists an exact sequence

$$(9) \quad 0 \rightarrow C_r^*(Y, \mathcal{G}) \rightarrow \mathcal{E} \rightarrow \mathcal{C}^\infty(S_{\text{vert}}^* Y) \rightarrow 0, \quad \mathcal{E} := \psi_{\text{inv}}^0(Y) + C_r^*(Y, \mathcal{G}),$$

obtained using the results of [18]. The operator A (or, rather, the family of operators $A = (A_b)$) has an invertible principal symbol, and hence the family $T = (T_b)$,

$$T_b := (1 + A_b^* A_b)^{-1/2} A_b,$$

consists of elliptic, $\mathcal{G}$ -invariant operators. Moreover, the operator T is an element of $\mathcal{E} := \psi_{\text{inv}}^0(Y) + C_r^*(Y, \mathcal{G})$, its principal symbol is invertible, and hence it defines a class $[T] \in K_1(\mathcal{C}^\infty(S_{\text{vert}}^* Y)) \simeq K^1(S_{\text{vert}}^* Y)$. Let

$$\partial : K_1^{\text{alg}}(S_{\text{vert}}^* Y) \rightarrow K_0^{\text{alg}}(C_r^*(Y, \mathcal{G})) \simeq K_0(C_r^*(Y, \mathcal{G}))$$

be the boundary map in the K -theory exact sequence

$$\begin{aligned} K_1^{\text{alg}}(C_r^*(Y, \mathcal{G})) &\rightarrow K_1^{\text{alg}}(\mathcal{E}) \rightarrow K_1^{\text{alg}}(S_{\text{vert}}^* Y) \xrightarrow{\partial} K_0^{\text{alg}}(C_r^*(Y, \mathcal{G})) \\ &\rightarrow K_0^{\text{alg}}(\mathcal{E}) \rightarrow K_0^{\text{alg}}(S_{\text{vert}}^* Y) \end{aligned}$$

associated to the exact sequence (9). Because $K_0(C_r^*(Y, \mathcal{G})) \simeq K_0(C_r^*(\mathcal{G}))$, by Lemma 2, we get a group morphism

$$(10) \quad \text{ind}_{\mathcal{G}} : K_1^{\text{alg}}(\mathcal{C}^\infty(S_{\text{vert}}^* Y)) \rightarrow K_0(C_r^*(\mathcal{G})),$$

which we shall call *the gauge-equivariant index morphism*. The image of A under the composition of the above maps is $\text{ind}_{\mathcal{G}}([T])$. We shall also write $\text{ind}_{\mathcal{G}}(A) :=$

$\text{ind}_{\mathcal{G}}([T])$, and call $\text{ind}_{\mathcal{G}}(A)$ the *gauge-equivariant index* of A . A more direct but longer definition is contained in the proof of Theorem 1 below (see Equation (13)). The gauge-equivariant index morphism descends in this case to a group morphism $K_1^{\text{alg}}(\mathcal{C}^\infty(S_{\text{vert}}^* Y)) \rightarrow K_0(C_r^*(\mathcal{G}))$ denoted in the same way. If $\mathcal{G}$ is a vector bundle, we can replace $\mathcal{E}$ with $\Psi_{\text{inv}}^0(Y)$ and $C_r^*(\mathcal{G})$ with $\Psi_{\text{inv}}^{-1}(Y)$.

We denote by I_N the unit of the matrix algebra $M_N(\mathcal{E})$. If $A \in \psi_{\text{inv}}^0(Y; E)$, we can find N large such that $\psi_{\text{inv}}^0(Y; E) \subset M_N(\psi_{\text{inv}}^0(Y))$ with 1 mapping to the projector e under this isomorphism. Then $A + (I_N - e)$ is invertible in this matrix algebra, and we define then $\text{ind}_{\mathcal{G}}(A) = \text{ind}_{\mathcal{G}}(A + I_N - e)$.

Let us briefly see what needs to be changed when $Y/\mathcal{G}$ is not compact. If $Y/\mathcal{G}$ is not compact, the above definition of the gauge-equivariant index applies only to elliptic operators in $I_N + M_N(\psi_{\text{inv}}^0(Y; E))$. However, all results below extend to operators in $I_N + M_N(\psi_{\text{inv}}^0(Y; E))$, after some obvious changes. For differential operators acting between sections of *different* bundles, we can define the gauge-equivariant index using the adiabatic groupoid of $\mathcal{G}$ as in [18]. For arbitrary elliptic pseudodifferential operators $A \in \psi_{\text{inv}}^m(Y; E_0, E_1)$, acting between sections of possibly different vector bundles, we can define the gauge-equivariant index using Kasparov's bivariant K -Theory. Since, when $Y/\mathcal{G}$ is non-compact, no element in $\psi_{\text{inv}}^m(Y; E_0, E_1)$ is invertible modulo regularizing operators, we must allow more general operators in this case. For example, we can take

$$A \in \Gamma(Y, \text{Hom}(E_0, E_1))^{\mathcal{G}} + \psi_{\text{inv}}^0(Y; E_0, E_1),$$

provided that it is bounded. If $A \in \psi_{\text{inv}}^m(Y; E_0, E_1)$, $m > 0$, we replace A by $(1 + AA^*)^{-1/2}A$, which will be an operator in the closure of $\psi_{\text{inv}}^0(Y; E_0, E_1)$, by the results of [21]. If A is elliptic, then the resulting family provides directly from the definition an element

$$(11) \quad \text{ind}_{\mathcal{G}}(A) \in KK(\mathbb{C}, C_r^*(\mathcal{G})).$$

To obtain the gauge-equivariant index as an element of $K_0(C_r^*(\mathcal{G}))$, we use the isomorphism $KK(\mathbb{C}, C_r^*(\mathcal{G})) \cong K_0(C_r^*(\mathcal{G}))$. We can also work with $\Psi_{\text{inv}}^{-\infty}(\mathcal{G})$ instead of $C_r^*(\mathcal{G})$, by using the smooth versions of KK -theory introduced in [33, 34].

The main property of the gauge-equivariant index of an operator A is that it gives the obstruction to the existence of invertible perturbations of A by lower order operators. We denote by $H^s(Y_b)$ the s th Sobolev space of $1/2$ -densities on Y_b , which is uniquely defined because of the bounded geometry of Y_b , for $Y/\mathcal{G}$ compact. More precisely, $H^s(Y_b)$ is, by definition, the domain of $(1 + D^*D)^{s/2m}$, if $D \in \psi_{\text{inv}}^m(Y)$ is elliptic and $s \geq 0$. For $s \leq 0$, $H^s(Y_b)$ is, by definition, the dual of $H^{-s}(Y_b)$. If $E \rightarrow Y$ is a $\mathcal{G}$ -equivariant vector bundle, then we shall denote by $H^s(Y_b, E)$ the Sobolev space of sections of E , defined analogously.

Theorem 1. *Let $\mathcal{G} \rightarrow B$ be a bundle of Lie groups acting on the fiber bundle $Y \rightarrow B$, as above, and assume that $Y/\mathcal{G}$ is compact, of positive dimension. Let $A \in \psi_{\text{inv}}^m(Y, E_0, E_1)$ be an elliptic operator. Then we can find $R \in \psi_{\text{inv}}^{m-1}(Y, E_0, E_1)$ such that*

$$A_b + R_b : H^s(Y_b, E_0) \rightarrow H^{s-m}(Y_b, E_1)$$

is invertible for all $b \in B$ if, and only if, $\text{ind}_{\mathcal{G}}(A) = 0$. Moreover, if $\text{ind}_{\mathcal{G}}(A) = 0$, then we can choose $R \in \psi_{\text{inv}}^{-\infty}(Y)$. The same result holds if $Y/\mathcal{G}$ is non-compact and $A \in \Gamma(Y, \text{Hom}(E_0, E_1))^{\mathcal{G}} + \psi_{\text{inv}}^0(Y; E_0, E_1)$ is elliptic and bounded.

Proof. We shall assume that $E_0 = E_1$ are trivial vector bundles of rank n , for simplicity of notation. The proof in the general case is the same. It is clear from definition that if we can find R with the desired properties, then $\text{ind}_{\mathcal{G}}(A) = 0 \in K^0(\mathfrak{g}^*)$. Suppose now that $A \in \psi_{\text{inv}}^m(Y, \mathbb{C}^N)$ is elliptic and has vanishing gauge-equivariant index. Using the notation $T = (1 + AA^*)^{-1/2}A$, we see that A_b is invertible between the indicated Sobolev spaces if, and only if, T_b is invertible as a bounded operator on $L^2(Y_b)^N$. We can hence assume that $m = 0$ and $T = A$.

Because $C_r^*(Y, \mathcal{G})$ satisfies $C_r^*(Y, \mathcal{G}) \simeq C_r^*(\mathcal{G}) \widehat{\otimes} \mathcal{K}$, by Lemma 2, we can use some general techniques to prove that the vanishing of $\text{ind}_{\mathcal{G}}(A)$ implies that A has a perturbation by $\mathcal{G}$ -invariant, regularizing operators in $\psi_{\text{inv}}^{-\infty}(Y, \mathbb{C}^N)$ that is invertible on each fiber. We fix an isomorphism $M_N(C_r^*(Y, \mathcal{G})) \cong C_r^*(\mathcal{G}) \otimes \mathcal{K}$. We now review this general technique using a generalization of an argument from [33]. Let $\mathcal{E}$ be the algebra introduced in Equation (9). We denote by I_N the unit of the matrix algebra $M_N(\mathcal{E})$. Also, denote by $\overline{\mathcal{E}}$ the closure of $\mathcal{E}$ in norm.

Choose a sequence of projections $p_n \in \mathcal{K}$, $\dim p_n = n$, such that $p_n \rightarrow 1$ in the strong topology. Because A_b is invertible modulo $C_r^*(\mathcal{G}_b) \otimes \mathcal{K}$, we can find a large n and $R \in M_N(\psi_{\text{inv}}^{-\infty}(Y))$ such that

$$A'_b := A_b \oplus R_b : L^2(Y_b)^N \oplus L^2(Y_b)^N \rightarrow L^2(Y_b)^N$$

is surjective and $(1 \otimes p_n)R_b(1 \otimes p_n) = R_b$, for all $b \in B$. Then $\{0\}$ is not in the spectrum of $A'A'^*$, and we can consider $V := (A'A'^*)^{-1/2}A' \in M_N(\overline{\mathcal{E}})$, which by construction satisfies $VV^* = I_N \in M_N(\overline{\mathcal{E}})$. Consequently, V^*V is a projection in $M_{2N}(\overline{\mathcal{E}})$. Because A is invertible modulo $M_N(C_r^*(\mathcal{G}) \otimes \mathcal{K})$,

$$V^*V - VV^* \in M_{2N}(C_r^*(\mathcal{G}) \otimes \mathcal{K}).$$

Let $e = I_N \oplus (1 \otimes p_n) - V^*V$, which is also a projection, by construction. Moreover,

$$(12) \quad e - (1 \otimes p_n) \in M_{2N}(C_r^*(\mathcal{G}) \otimes \mathcal{K}),$$

and hence both e and $1 \otimes p_n$ are projection in $M_{2N}(C_r^*(\mathcal{G})^+ \otimes \mathcal{K})$ (for any algebra B , we denote by B^+ the algebra with adjoint unit). Equation (12) gives that, by definition, $[e] - [1 \otimes p_n]$ defines a K -theory class in $K_0(C_r^*(\mathcal{G}))$. From definition, we get then

$$(13) \quad \text{ind}_{\mathcal{G}}(A) = [e] - [1 \otimes p_n].$$

Now, if $\text{ind}_{\mathcal{G}}(A) = 0$, then we can find k such that $e \oplus 1 \otimes p_k$ is (Murray-von Neumann) equivalent to $1 \otimes p_{n+k}$. By replacing our original choice for n with $n+k$, we may assume that e and $1 \otimes p_n$ are equivalent, and hence that we can find $u \in \mathbb{C}I_{2N} + M_{2N}(C_r^*(\mathcal{G}) \otimes \mathcal{K})$ with the following properties: there exists a large l and $x \in M_{2N}(C_r^*(\mathcal{G}))$ such that, if we denote $e_0 = 1 \otimes p_l \oplus 1 \otimes p_n$, then $u = I_{2N} + e_0 x e_0$ and $e = u(1 \otimes p_n)u^{-1}$. Then Vu is in $M_n(\overline{\mathcal{E}})$ (more precisely $I_N V I_N = V$) and is invertible. Consequently $B := (A'A'^*)^{1/2}V$ is also invertible. But B is a perturbation of A' , and hence also of A , by an element in $M_{2N}(C_r^*(\mathcal{G}) \otimes \mathcal{K})$. Since $\psi_{\text{inv}}^{-\infty}(Y)$ is dense in $C_r^*(\mathcal{G})$, this gives the result. $\square$

4. THE GAUGE-EQUIVARIANT INDEX: A FORMULA

We now treat in more detail the case of bundles of solvable Lie groups, when more precise results can be obtained. Other classes of groups will lead to completely different problems and results, so we leave their study for the future. The class of simply-connected solvable fibers is rich enough and has many specific features, so

we content ourselves from now on with this case only.

Assumption. *From now on and throughout this paper, we shall assume that the family $\mathcal{G}$ consists of simply-connected solvable Lie groups. By “simply-connected” we mean, as usual, “connected with trivial fundamental group.”*

We shall denote by $\mathfrak{g} \rightarrow B$ the bundle of Lie algebras of the groups $\mathcal{G}_b$, $\mathfrak{g}_b \simeq \text{Lie } \mathcal{G}_b$ and by

$$\exp : \mathfrak{g} \rightarrow \mathcal{G}.$$

the exponential map.

In order to study the algebra $C_r^*(\mathcal{G})$ and its K -groups, we shall deform it to a commutative algebra. This deformation is obtained as follows. Let $\mathcal{G}_{adb} = \{0\} \times \mathfrak{g} \cup (0, 1] \times \mathcal{G}$, $B_1 = [0, 1] \times B$, and $p_{adb} : \mathcal{G}_{adb} \rightarrow B_1$ be the natural projection. On $\mathcal{G}_{adb}$ we put the smooth structure induced by

$$\phi : B_1 \times \mathfrak{g} \rightarrow \mathcal{G}_{adb}$$

$\phi(0, X) = (0, X)$ and $\phi(t, X) = (t, \exp(tY))$, which is a bijection for all $(t, X) \in [0, 1] \times \mathfrak{g}$ in a small neighborhood of $\{0\} \times \mathfrak{g} \cup B_1 \times \{0\}$. Then we endow $\mathcal{G}_{adb}$ with the Lie bundle structure induced by the pointwise product. Evaluation at $t \in [0, 1]$ induces algebra morphisms

$$e_t : \psi_{\text{inv}}^m(\mathcal{G}_{adb}) \rightarrow \psi_{\text{inv}}^m(\mathcal{G}), \quad t > 0,$$

and

$$e_0 : \psi_{\text{inv}}^m(\mathcal{G}_{adb}) \rightarrow \psi_{\text{inv}}^m(\mathfrak{g}), \quad t = 0.$$

Passing to completions, we obtain morphisms e_t from $C_r^*(\mathcal{G}_{adb})$ to $C_r^*(\mathcal{G})$, for $t > 0$, and to $C_r^*(d^{-1}(0) \times B) \simeq C_0(\mathfrak{g}^*)$, for $t = 0$.

Lemma 3. *The morphisms $e_t : C_r^*(\mathcal{G}_{adb}) \rightarrow C_r^*(\mathcal{G})$, for $t > 0$, and $e_0 : C_r^*(\mathcal{G}_{adb}) \rightarrow C_0(\mathfrak{g}^*)$, for $t = 0$, induce isomorphisms in K -theory.*

Proof. Assume first that there exists a Lie group bundle morphism $\mathcal{G} \rightarrow B \times \mathbb{R}$. (In other words, there exists a smooth map $\mathcal{G} \rightarrow \mathbb{R}$ that is a morphism on each fiber.) Let $\mathcal{G}'$ denote the kernel of this morphism and let $\mathcal{G}'_{adb}$ be obtained from $\mathcal{G}'$ by the same deformation construction by which $\mathcal{G}_{adb}$ was obtained from $\mathcal{G}$. Then we obtain a smooth map $\mathcal{G}_{adb} \rightarrow \mathbb{R}$ that is a group morphism on each fiber, and hence

$$C_r^*(\mathcal{G}_{adb}) \simeq C_r^*(\mathcal{G}'_{adb}) \rtimes \mathbb{R}, \quad C_r^*(\mathcal{G}) \simeq C_r^*(\mathcal{G}') \rtimes \mathbb{R}, \quad \text{and} \quad C_0(\mathfrak{g}^*) \simeq C_0(\mathfrak{g}'^*) \rtimes \mathbb{R}.$$

Moreover, all above isomorphisms are natural, and hence compatible with the morphisms e_t . Assuming now that the result was proved for all Lie group bundles of smaller dimension, we obtain the desired result for $\mathcal{G}$ using Connes' Thom isomorphism in K -theory [13], which in this particular case gives:

$$K_i(C_r^*(\mathcal{G}_{adb})) \simeq K_{i+1}(C_r^*(\mathcal{G}'_{adb})), \quad K_i(C_r^*(\mathcal{G})) \simeq K_{i+1}(C_r^*(\mathcal{G}')), \quad \text{and} \\ K_i(C_0(\mathfrak{g}^*)) \simeq K_{i+1}(C_0(\mathfrak{g}'^*)).$$

This will allow us to complete the result in the following way. Let U_k be the open subset of B consisting of those $b \in B$ such that $[\mathcal{G}_b, \mathcal{G}_b]$ has dimension $\geq k$. From the Five Lemma and the six term exact sequences in K -theory associated to the ideal $C_r^*(\mathcal{G}_{adb}|_{U_k \setminus U_{k+1}})$ of $C_r^*(\mathcal{G}_{adb}|_{U_{k-1} \setminus U_{k+1}})$, for each k , we see that it is enough to prove our result for $\mathcal{G}_{adb}|_{U_k \setminus U_{k+1}}$ for all k . Thus, by replacing B with

$U_k \setminus U_{k+1}$, we may assume that the rank of the abelianization of $\mathcal{G}_b$ is independent of b . Consequently, the abelianizations of $\mathcal{G}_b$ form a vector bundle

$$\mathcal{A} := \cup \mathcal{G}_b / [\mathcal{G}_b, \mathcal{G}_b]$$

on B .

A similar argument, using the Meyer-Vietoris exact sequence in K -theory and the compatibility of K -theory with inductive limits [12], shows that we may also assume the vector bundle $\mathcal{A}$ of abelianizations to be trivial. Then the argument at the beginning of the proof applies, and the result is proved. $\square$

From the above lemma we immediately obtain the following corollary:

Corollary 1. *Let $\mathcal{G}$ be a bundle of simply connected, solvable Lie groups. Then*

$$K_i(C_r^*(\mathcal{G})) \simeq K_i(C_0(\mathfrak{g}^*)) \simeq K^i(\mathfrak{g}^*).$$

We now give an interpretation of $\text{ind}_{\mathcal{G}}(A)$, for $\mathcal{G}$ a vector bundle, using the properties of the indicial family $\hat{A}(\tau)$ of A . We assume that $Y/\mathcal{G}$ is compact.

We shall also use the following construction. Let X be a compact manifold with boundary. Let $T(x)$ be a family of elliptic pseudodifferential operators acting between sections of two smooth vector bundles, E_0 and E_1 , on the fibers of a fiber bundle $M \rightarrow X$ whose fibers are compact manifolds without corners. Then we can realize the index of T as an element in the relative group $K^0(X, \partial X)$. This can be done directly using Kasparov's theory or by the "Atiyah-Singer trick" as follows. We proceed as in [8], Proposition (2.2), to define a smooth family of maps $R(x) : \mathbb{C}^N \rightarrow \mathcal{C}^\infty(Y)$, such that the induced map

$$V := T \oplus R : \mathcal{C}^\infty(X)^N \oplus \mathcal{C}^\infty(X, L^2(Y; E_0)) \rightarrow \mathcal{C}^\infty(X, L^2(Y; E_1))$$

is onto for each x . Since $T(x)$ is invertible for $x \in \partial X$, we can choose $R(x) = 0$ for $x \in \partial X$. Then $\ker(V)$ is a vector bundle on X , which is canonically trivial on the boundary ∂X . The general definition of the index of the family T in [8] is that of the difference of the kernel bundle $\ker(V)$ and the trivial bundle $X \times \mathbb{C}^N$. Since the bundle $\ker(V)$ is canonically trivial on the boundary of X , we obtain an element

$$(14) \quad \deg(T) \in K^0(X, \partial X).$$

The degree is invariant with respect to homotopies T_t of families of operators on X that are *invertible* on ∂X throughout the homotopy. We shall use the degree in Theorem 3 for $X = B_R$, a large closed ball in $\mathcal{G}^* \simeq \mathfrak{g}^*$, or for X being the radial compactification of $\mathfrak{g}^*$. If the boundary of X is empty, this construction goes back to Atiyah and Singer and then $\deg(T)$ is simply the family index of T . This definition of $\deg(T)$ is due to Melrose. We note that when $\partial X \neq \emptyset$, the degree is not a local quantity in T , in the sense that it depends on more than just the principal symbol.

Assume now that the family T above consists of order zero operators and $T(x)$ is a multiplication operator for each $x \in \partial X$. We want to compute the Chern character of $\deg(T)$ using the Atiyah-Singer family index formula [7, 8]. To introduce the main ingredients of the index formula, denote by $S_{vert}^* M$ the set of unit vectors in the dual of the vertical tangent bundle $T_{vert} M$ to the fibers of $M \rightarrow X$. Because the family T is elliptic, the principal symbols define an invertible matrix of functions

$$a = \sigma_0(T) \in \mathcal{C}^\infty(S_{vert}^* M; \text{Hom}(E_0, E_1)).$$

Since the operators $T(x)$ are multiplication operators, we can then extend a to an invertible endomorphism on

$$S_M := S_{vert}^* M \cup B^*,$$

with B^* denoting the set of vertical cotangent vectors of length ≤ 1 above ∂X , as in [5]. The constructions of [7, 8] are in terms of $[a'] \in K^0(T_{vert}^* M, T_{vert}^* M|_{\partial M})$ obtained by applying the clutching (or difference) construction to a . Explicitly, $[a']$ is represented by (E_0, E_1, a_1) (where a_1 is a smooth function that coincides with a outside a neighborhood of the zero section). It defines an element in

$$K^0(T_{vert}^* M, T_{vert}^* M|_{\partial M}) = K^0(T_{vert}^* M \setminus T_{vert}^* M|_{\partial M}),$$

because a defines an endomorphism of the trivial bundle $\mathbb{C}^N$ which is invertible outside a compact set. (Recall from [3] that we can represent the K -theory groups of a space X as equivalence classes of triples (E_0, E_1, a) , where E_0 and E_1 are vector bundles on X and a is a vector bundle morphism $E_0 \rightarrow E_1$ which is an isomorphism outside a compact set).

When $E_0 \cong E_1$, we can assume that $E_0 = E_1 = M \times \mathbb{C}^N$ are trivial of rank N , and we have that a is an invertible matrix valued function, which hence defines an element $[a] \in K^1(S_M)$. Let B_1 be the set of vectors of norm at most 1 in $T^* M$. After the identification (up to homeomorphism) of $B_1 \setminus S_M$, the interior of S_M , with the difference $T_{vert}^* M \setminus T_{vert}^* M|_{\partial M}$, we have

$$[a'] = \partial[a].$$

We denote by $\pi_* : H^*(S_M) \rightarrow H^{*-2n+1}(X, \partial X)$ the integration along the fibers, where n is the dimension of the fibers of $M \rightarrow X$. Integration along the fibers in this case is the composition of

$$\partial : H^*(S_M) \rightarrow H^{*+1}(B_1, S_M) \cong H_c^{*+1}(T_{vert}^* M \setminus T_{vert}^* M|_{\partial M})$$

and

$$\tilde{\pi}_* : H_c^*(T_{vert}^* M \setminus T_{vert}^* M|_{\partial M}) \rightarrow H_c^{*-2n}(X \setminus \partial X)$$

obtained by integration along the fibers of the bundle $T_{vert}^* M \setminus T_{vert}^* M|_{\partial M} \rightarrow X \setminus \partial X$:

$$\pi_* = \tilde{\pi}_* \circ \partial.$$

To state the following result, we also need $Ch : K_1(S_M) \rightarrow H^{odd}(S_M)$, the Chern character in K -Theory and $\mathcal{T}$, the Todd class of $(T_{vert}^* M) \otimes \mathbb{C}$, the complexified vertical tangent bundle of the fibration $M \rightarrow X$, as in [8]. Using the notation introduced above, we have:

Theorem 2. *Let $M \rightarrow X$ be a smooth fiber bundle whose fibers are smooth manifolds (without corners), and let T be a family of order zero elliptic pseudodifferential operators acting along the fibers of $M \rightarrow X$. Assume X is a manifold with boundary ∂X such that the operators $T(x)$ are multiplication operators on ∂X , also let $[a']$ and $[a]$ be the classes defined above. Then*

$$Ch(\deg(T)) = (-1)^n \tilde{\pi}_*(Ch[a']\mathcal{T}) \in H^*(X, \partial X),$$

If $E_0 \cong E_1$, then we also have $Ch(\deg(T)) = (-1)^n \pi_(Ch[a]\mathcal{T})$.*

Proof. For continuous families $T(x)$ that are multiplication operators on the boundary ∂X , the degree is a local quantity – it depends only on the principal symbol – so we can follow word for word [8] to prove that

$$Ch(\deg(T)) = (-1)^n \tilde{\pi}_*(Ch[a']\mathcal{T}) \in H^*(X, \partial X).$$

When $E_0 \cong E_1$, using $Ch[a'] = Ch(\partial[a]) = \partial Ch[a]$, we get

$$\begin{aligned} Ch(\deg(T)) &= (-1)^n \tilde{\pi}_*(\partial Ch[a]\mathcal{T}) = (-1)^n \tilde{\pi}_* \circ \partial(Ch[a]\mathcal{T}) \\ &= (-1)^n \pi_*(Ch[a]\mathcal{T}) \in H^*(X, \partial X). \end{aligned}$$

This completes the proof. $\square$

It is interesting to note that it is not possible in general to give a formula for $Ch(\deg(T))$ only in terms of its principal symbol, without further assumptions on T (for example that T consists of multiplication operators on the boundary, as in our theorem). A consequence is that, in general, the formula for $Ch(\deg(T))$ will involve some non-local invariants. It would be nevertheless useful to find such a formula.

Returning to our considerations, we continue to assume that $Z := Y/\mathcal{G}$ is compact, and we fix a metric on $\mathcal{G}$ (which, we recall, is a vector bundle in these considerations). If $A \in \Psi_{\text{inv}}^\infty(Y; E_0, E_1)$ is elliptic (in the sense that its principal symbol is invertible outside the zero section), then the indicial operators $\hat{A}(\tau)$ are invertible for $|\tau| \geq R$, $\tau \in \mathcal{G}^*$, and some large R . In particular, by restricting the family $\hat{A}$ to the ball

$$B_R := \{|\tau| \leq R\},$$

we obtain a family of elliptic operators that are invertible on the boundary of B_R , and hence $\hat{A}$ defines an element

$$(15) \quad \deg_{\mathcal{G}}(A) := \deg(\hat{A}) \in K^0(B_R, \partial B_R) \simeq K^0(\mathfrak{g}^*)$$

in the K -group of the ball of radius R , relative to its boundary, as explained above, called also the *degree* of A .

We want a formula for the Chern character of the degree of $A \in \Psi_{\text{inv}}^\infty(Y; E_0, E_1)$. Because A is elliptic, its principal symbol defines a class $[a'] \in K^0((T_{\text{vert}}^*Y)/\mathcal{G})$. If E_0 and E_1 are isomorphic, then it also defines a class $[a] \in K^0((S_{\text{vert}}^*Y)/\mathcal{G})$. Denote by n the dimension of the quotient $Z_b = Y_b/\mathcal{G}_b$ (which is independent of b because we assumed B connected), and let $\pi : (S_{\text{vert}}^*Y)/\mathcal{G} \rightarrow B$ be the projection and

$$(16) \quad \begin{aligned} \tilde{\pi}_* : H^*((T_{\text{vert}}^*Y)/\mathcal{G}) &\rightarrow H_c^{*-2n}(\mathfrak{g}^*) \\ \pi_* : H^*((S_{\text{vert}}^*Y)/\mathcal{G}) &\rightarrow H_c^{*-2n+1}(\mathfrak{g}^*) \end{aligned}$$

be the integration along the fibers in cohomology. Then

$$Ch(\deg_{\mathcal{G}}(A)) \in H_c^*(\mathfrak{g}^*) \simeq H^{*+n}(B, \mathcal{O}),$$

and the following theorem gives a formula for this cohomology class in terms of the classes $[a']$ or $[a]$ defined above.

We denote by $\mathcal{T}$ the Todd class of the vector bundle $(T_{\text{vert}}Y)/\mathcal{G} \otimes \mathbb{C} \rightarrow Y/\mathcal{G}$. We assume B to be compact.

Theorem 3. *If $A \in \Psi_{\text{inv}}^\infty(Y; E_0, E_1)$ is elliptic, then the Chern character of $\deg_{\mathcal{G}}(A)$ is given by*

$$Ch(\deg_{\mathcal{G}}(A)) = (-1)^n \tilde{\pi}_*(Ch[a']\mathcal{T}) \in H_c^*(\mathfrak{g}^*),$$

Moreover, $Ch(\deg_{\mathcal{G}}(A)) = (-1)^n \pi_*(Ch[a]\mathcal{T})$ if $E_0 \cong E_1$.

Observations. It is almost always the case that $E_0 \cong E_1$. For example, it is easy to see that this must happen if the Euler characteristic of $(T_{\text{vert}}^*Y)/\mathcal{G}$ vanishes. This assumption is satisfied if $\mathcal{G} = B \times \mathbb{R}^q$, $q > 0$, which is the case for parameter dependent pseudodifferential operators, for example.

Another observation is that if the set of elliptic elements in $\Psi_{\text{inv}}^\infty(Y; E_0, E_1)$ is not empty, then $Z := Y/\mathcal{G}$ is compact.

Proof. We cannot use Theorem 2 directly because our family $\hat{A}$ does not consist of multiplication operators on the boundary. Nevertheless, we can deform $\hat{A}$ to a family of operators that are multiplication operators at ∞ , for suitable A . We now construct this deformation.

Let E be a vector bundle over V . We consider classical symbols $S_c^m(E)$ whose support projects onto a compact subset of V . Let

$$A_Y := (T_{\text{vert}}Y)/\mathcal{G} \cong T_{\text{vert}}Z \times_B \mathfrak{g}.$$

First we need to define a nice quantization map $q : S^m(A_Y^*) \rightarrow \psi_{\text{inv}}^m(Y)$. To this end, we proceed as usual, using local coordinates, local quantization maps, and partitions of unity, but being careful to keep into account the extra structure afforded by our settings: the fibration over B and the action of $\mathcal{G}$. Here are the details of how this is done.

Fix a cross section for $Y \rightarrow Z := Y/\mathcal{G}$ and, using it, identify Y with $Z \times_B \mathcal{G}$ as $\mathcal{G}$ -spaces. Denote by $p_0 : Z \rightarrow B$ the natural projection. We cover B with open sets $U_{\alpha'}$ that are diffeomorphic to open balls in a Euclidean space such that $Z|_{U_{\alpha'}} \cong U_{\alpha'} \times F$ and $\mathcal{G}|_{U_{\alpha'}} \cong U_{\alpha'} \times \mathbb{R}^q$. We also cover F with open domains of coordinate charts $V_{\alpha''}$. Then we let $W_\alpha = p_0^{-1}(U_{\alpha'}) \cap V_{\alpha''}$, with $\alpha = (\alpha', \alpha'')$. The natural coordinate maps on W_α then give rise to a quantization map

$$(17) \quad q_\alpha : S_c^m(A_Y^*|_{W_\alpha}) \rightarrow \psi_{\text{inv}}^m(W_\alpha \times_B \mathcal{G}), \quad q_\alpha(a) = a(b, x, D_x, D_t),$$

where we identify

$$(18) \quad S_c^m(A_Y^*|_{W_\alpha}) = S_c^m(T_{\text{vert}}^*(W_\alpha) \times_B \mathcal{G}) = S_c^m(U_{\alpha'} \times T^*V_{\alpha''} \times \mathbb{R}^{*q}).$$

Denote by $b \in U_{\alpha'}$, $(x, y) \in T^*V_{\alpha''} \cong V_{\alpha''} \times \mathbb{R}^{*n}$, and $\tau \in \mathbb{R}^{*q}$ the corresponding coordinate maps. Then $q_\alpha(a) = a(b, x, D_x, D_t)$ acts on $\mathcal{C}_c^\infty(U_{\alpha'} \times T^*V_{\alpha''} \times \mathbb{R}^q)$ as

$$\begin{aligned} & a(b, x, D_x, D_t)u(b, x, t) \\ &= (2\pi i)^{-n-q} \int_{\mathbb{R}^{n+q}} \left(\int_{\mathbb{R}^{n+q}} e^{i(x-z) \cdot y + i(t-s) \cdot \tau} a(b, x, y, \tau) u(b, z, s) ds dz \right) dy d\tau. \end{aligned}$$

Choose now a partition of unity ϕ_α^2 subordinated to W_α and let

$$q(a) = \sum_{\alpha} q_\alpha(\phi_\alpha a) \phi_\alpha.$$

The main properties of q are the following:

1. if a has order m , then $\sigma_m(q(a)) = a$, modulo symbols of lower order;

2. there exist maps $q_b : S^m(T^*Z_b) \rightarrow \Psi^m(Z_b)$ such that

$$\widehat{q(a)}(\tau) = q_b(a(\cdot, \tau)) = a(b, x, \tau + D_x),$$

if $\tau \in \mathfrak{g}_b^*$ and $x \in Z_b$;

3. the maps q_b define a quantization map

$$\tilde{q} : \Gamma(\mathfrak{g}^*, S^m(T_{vert}^*Z)) \rightarrow \Gamma(\mathfrak{g}^*, \Psi^m(Z)),$$

where we regard $\Psi^m(Z_b)$ as defining a bundle of algebras, $\Psi^m(Z)$, on B , first, and then on $\mathfrak{g}^*$, by pull-back. Similarly, we regard $S^m(T_{vert}^*Z)$ as defining a bundle over Z , which we then pull back to a bundle on $\mathfrak{g}^*$. (See also the discussion related to Equations (7) and (8).)

The deformation of our family is obtained as follows. Let $|\tau|$ and $|y|$ be the norms $\tau \in \mathfrak{g}^*$ and $y \in T_{vert}^*Z$. Define then

$$(19) \quad \phi_\lambda^2(y, \tau) = 1 + \lambda|y|^2 + \lambda(1 + \lambda|\tau|^2)^{-1}|y|^2 \quad \text{and} \quad \psi_\lambda^{-2}(y, \tau) = 1 + \lambda|\tau|^2,$$

which were chosen to satisfy

$$1 + \phi_\lambda^2|\tau|^2 + \psi_\lambda^2|y|^2 = 1 + |\tau|^2 + |y|^2 + \lambda|\tau|^2|y|^2.$$

For any symbol $a \in S^0(A_Y^*)$, $A_Y^* = T_{vert}^*Z \times_B \mathfrak{g}^*$, we let

$$a_{\lambda, \tau}(y) = a(\psi_\lambda y, \phi_\lambda \tau), \quad \lambda \in [0, 1], \quad \tau \in \mathfrak{g}_b^*, \quad \text{and} \quad y \in T^*Z_b.$$

We can define then $A_\lambda(\tau) := q_b(a_{\lambda, \tau})$, $\tau \in \mathfrak{g}_b^*$, which is the same as saying that $A_\lambda = \tilde{q}(a_{\lambda, \tau})$, and these operators will satisfy the following properties:

1. For each fixed λ , the operators $A_\lambda(\tau)$ define a section of $\Psi^m(Z)$ over $\mathfrak{g}^*$ and these sections depend smoothly on λ (in other words, $A_\lambda(\tau)$ depends smoothly on both λ and τ , in any trivialization);
2. $A_0(\tau) = \widehat{q(a)}(\tau)$, for all τ ;
3. For each nonzero $\tau' \in \mathfrak{g}^*$ and $\lambda > 0$, the limit $\lim_{t \rightarrow \infty} A_\lambda(t\tau')$ exists and is a multiplication operator;
4. If $a, b \in S^0(A_Y^*)$ are such that $ab = 1$, a is homogeneous of order zero outside the unit ball, and if we define $A_\lambda := \tilde{q}(a_{\lambda, \tau})$ and $B_{\lambda, \tau} := \tilde{q}(b_{\lambda, \tau})$, then there exists a constant $C > 0$ such that

$$\|A_\lambda(\tau)B_{\lambda, \tau} - 1\| \leq C(1 + |\tau|)^{-1}$$

and similarly

$$\|B_{\lambda, \tau}A_\lambda(\tau) - 1\| \leq C(1 + |\tau|)^{-1}$$

For all τ and λ ;

5. All these estimates extend in an obvious way to matrix valued symbols.

These properties are proved as follows. We first recall that, for any vector bundle E , we can identify the space of classical symbols $S_c^0(E)$ with $\mathcal{C}_c^\infty(E_1)$, the space of compactly supported functions on E_1 , the unit ball of E , by $E_1 \setminus \partial E_1 \ni \xi \rightarrow (1 - \|\xi\|^2)^{-1}\xi \in E$. For any quantization map, the norm of the resulting operator will depend on finitely many derivatives. Because we can extend $a_{\lambda, \tau}$ to a smooth function on the radial compactification of A_Y^* , the first property follows. The second property is obvious. The third property is obtained using the same argument and observing that, for $\lambda > 0$, we can further extend our function to the radial compactification in λ also. By investigating what this limit is along various rays, we obtain the third property.

The fourth property is obtained using the following observation: there exist a constant $C > 0$ and seminorms $\|\cdot\|_0$ on $S_c^0(T^*\mathbb{R}^n)$ and $\|\cdot\|_{-1}$ on $S^{-1}(T^*\mathbb{R}^n)$ such that, for any symbols $a, b \in S_c^0(T^*\mathbb{R}^n)$,

$$\|(ab)(x, D_x) - a(x, D_x)b(x, D_x)\| \leq \sum_j C(\|a\|_0 \|\partial_{y_j} b\|_{-1} + \|\partial_{y_j} a\|_{-1} \|b\|_0),$$

∂_{y_j} being all derivatives in the symbolic directions (whose coordinates are denoted by y). Finally, the fifth property is obvious.

We now turn to the proof of the formula for the degree of A stated in our theorem. We prove it by a sequence of successive reductions, using the facts established above. First, it is easy to see that $\deg_{\mathcal{G}}(A)$ depends only on its principal symbol, and hence we can assume that A has order zero and $A = q(a)$, where $a = \sigma_0(A)$.

The above deformation can be used to prove our theorem as follows. Fix R large enough, and restrict the families A_λ to the closed ball of radius R in $\mathfrak{g}^*$. For $|\tau| = R$ large enough, all operators $A_\lambda(\tau)$, $\lambda \in [0, 1]$ are invertible, so the degree $\deg(A_\lambda)$ of these families is defined and does not depend on λ or R , provided that R is large enough. Since $\deg_{\mathcal{G}}(A) = \deg(A_0)$, by definition, it is enough to compute $\deg(A_\lambda)$, for any given λ . Choose then $\lambda > 0$ arbitrary, and let $R \rightarrow \infty$. Then the family A_λ extends to a continuous family on the radial compactification of $\mathfrak{g}^*$, which consists of multiplication operators on the boundary. Moreover, the symbol class of A_λ is nothing by the extension of $a_\lambda(\tau)$ to the radial compactification in τ and λ (which is a manifold with corners of codimension two).

We can use then Theorem 2 to conclude that

$$\deg(A_\lambda) = (-1)^n \tilde{\pi}_*(Ch[a'_\lambda]T) \in H_c^*(\mathfrak{g}^*).$$

But a_λ is homotopic to a through symbols that are invertible outside a fixed compact set, so $[a'_\lambda] = [a']$. We get

$$\deg(A) = (-1)^n \tilde{\pi}_*(Ch[a']T) \in H_c^*(\mathfrak{g}^*).$$

To obtain the second form of our formula for $E_0 \cong E_1$, and thus finish the proof, we proceed as at the end of the proof of Theorem 2, using $Ch[a'] = \partial Ch[a]$. $\square$

To prove the following result, we shall use terminology from algebraic topology: if $I_k \subset A_k$ are two-sided ideal of some algebras A_0 and A_1 and $\phi : A_0 \rightarrow A_1$ is an algebra morphism, we say that ϕ induces a morphism of pairs $\phi : (A_0, I_0) \rightarrow (A_1, I_1)$ if, by definition, $\phi(I_0) \subset I_1$.

Theorem 4. *Let $\mathcal{G}$ be a vector bundle and $A \in \Psi_{\text{inv}}^m(Y; \mathbb{C}^N)$ be an elliptic element. Then*

$$\text{ind}_{\mathcal{G}}(A) = \deg_{\mathcal{G}}(A) \in K^0(\mathfrak{g}^*).$$

Proof. By embedding E in a trivial vector bundle, we can reduce to the case $E = B \times \mathbb{C}^N$. Let $B_R = \{|\tau| \leq R\} \subset \mathcal{G}^*$ be as above. The algebra

$$\mathfrak{A}_R := \mathcal{C}^\infty(B_R, \Psi^\infty(Y_b))$$

of $\mathcal{C}^\infty$ -families of pseudodifferential operators on B_R acting on fibers of $Y \times_B B_R \rightarrow B_R$, contains as an ideal $\mathfrak{I}_R = \mathcal{C}_0^\infty(B_R, \Psi^{-\infty}(Y_b))$, the space of families of smoothing operators that vanish to infinite order at the boundary of B_R . If A is an elliptic family, as in the statement of the lemma, and if R is large enough, then $\hat{A}$, the indicial family of A , defines by restriction an element of $M_N(\mathfrak{A}_R)$ that is invertible modulo $M_N(\mathfrak{I}_R)$.

Recall that the boundary map ∂_1 in algebraic K -theory associated to the ideal $\mathfrak{I}_R$ of the algebra $\mathfrak{A}_R$ gives $\partial_1[A] = \deg_{\mathcal{G}}(A)$, by definition. Also, the boundary map ∂_0 in algebraic K -theory associated to the ideal $\Psi_{\text{inv}}^{-\infty}(Y)$ of the algebra $\Psi_{\text{inv}}^{\infty}(Y)$ gives $\partial_0[A] = \text{ind}_{\mathcal{G}}(A)$. We want to prove that $\partial_1[A] = \partial_0[A]$. The desired equality will follow by a deformation argument, which involves constructing an algebra smoothly connecting the ideals $\mathfrak{I}_R$ and $\Psi_{\text{inv}}^{-\infty}(Y)$.

Consider inside $\mathcal{C}^{\infty}([0, R^{-1}], \Psi_{\text{inv}}^{-\infty}(Y))$ the subalgebra of families $T = (T_x)$ such that $\hat{T}_x(\tau) = 0$ for $|\tau| \geq x^{-1}$. (In other words, $T_x \in \mathfrak{I}_{x^{-1}}$, if $x \neq 0$, and T_0 is arbitrary.) Denote this subalgebra by $\mathfrak{I}_{R\infty}$. Also, let $\mathfrak{A}_{R\infty}$ be the set of families $A = (A_x)$, $x \in [0, R^{-1}]$, $A_x \in \mathfrak{A}_{x^{-1}}$, if $x \neq 0$, $A_0 \in \Psi_{\text{inv}}^{\infty}(Y)$ arbitrary such that the families $AT := (A_x T_x)$ and $TA := (T_x A_x)$ are in $\mathfrak{I}_{R\infty}$, for all families $T = (T_x) \in \mathfrak{I}_{R\infty}$.

It follows that $\mathfrak{I}_{R\infty}$ is a two-sided ideal in $\mathfrak{A}_{R\infty}$ and that the natural restrictions of operators to $x = R^{-1}$ and, respectively, to $x = 0$, give rise to morphisms of pairs

$$e_1 : (\mathfrak{A}_{R\infty}, \mathfrak{I}_{R\infty}) \rightarrow (\mathfrak{A}_R, \mathfrak{I}_R), \quad \text{and} \\ e_0 : (\mathfrak{A}_{R\infty}, \mathfrak{I}_{R\infty}) \rightarrow (\Psi_{\text{inv}}^{\infty}(Y), \Psi_{\text{inv}}^{-\infty}(Y)).$$

Moreover, the indicial family of the operator A gives rise, by restriction to larger and larger balls B_r , to an invertible element in $\mathfrak{A}_{R\infty}$, also denoted by A . Let ∂ be the boundary map in algebraic K -theory associated to the pair $(\mathfrak{A}_{R\infty}, \mathfrak{I}_{R\infty})$. Then $(e_0)_* \partial[A] = \partial_0[A]$ and $(e_1)_* \partial[A] = \partial_1[A]$. Since $(e_0)_* : K_0(\mathfrak{I}_{R\infty}) \rightarrow K_0(\Psi_{\text{inv}}^{-\infty}(Y))$ and $(e_1)_* : K_0(\mathfrak{I}_{R\infty}) \rightarrow K_0(\mathfrak{I}_R)$ are natural isomorphisms, our result follows. $\square$

We now drop the assumption above that $\mathcal{G}$ consist of abelian Lie groups, assuming instead that $\mathcal{G}$ consists of simply-connected solvable Lie groups, and want to compute the Chern character of the gauge-equivariant index $\text{ind}_{\mathcal{G}}(A)$, for an elliptic family $A \in \psi_{\text{inv}}^m(Y)$. One difficulty that we encounter is that the space on which the principal symbols are defined, that is $(S_{\text{vert}}^* Y)/\mathcal{G}$, is not orientable in general. (Recall that $S_{\text{vert}}^* Y$ is the space of vectors of length one of $T_{\text{vert}}^* Y$, the dual of the vertical tangent bundle $T_{\text{vert}} Y$ to the fibers of $Y \rightarrow B$.)

We denote by $\mathcal{T}$ the Todd class of the vector bundle $(T_{\text{vert}} Y)/\mathcal{G} \otimes \mathbb{C} \rightarrow Y/\mathcal{G}$ and by π_* the integration along the fibers of $(S_{\text{vert}}^* Y)/\mathcal{G} \rightarrow B$, as above. We assume B to be compact.

Theorem 5. *Let $\mathcal{G}$ be a bundle of Lie groups whose fibers are simply-connected, solvable Lie groups. Let $A \in \psi_{\text{inv}}^m(Y, E)$ be an elliptic, $\mathcal{G}$ -invariant family, and let $[\sigma_m(A)] \in K^1((S_{\text{vert}}^* Y)/\mathcal{G})$ be the class defined by the principal symbol $\sigma_m(A)$ of A . Then the Chern character of the gauge-equivariant index of A is given by*

$$\text{Ch}(\text{ind}_{\mathcal{G}}(A)) = (-1)^n \pi_* (\text{Ch}[\sigma_m(A)] \mathcal{T}) \in H_c^*(\mathfrak{g}^*),$$

where n is the dimension of the fibers of $(S_{\text{vert}}^* Y)/\mathcal{G} \rightarrow B$.

Proof. Note first that we can deform the bundle of Lie groups $\mathcal{G}$ to the bundle of commutative Lie groups $\mathfrak{g}$ as before, using $\mathcal{G}_{\text{adb}}$. Moreover, we can keep the principal symbol of A constant along this deformation. This shows that we may assume $\mathcal{G}$ to consist of commutative Lie groups, i.e. that $\mathcal{G}$ is a vector bundle. The result then follows from Theorems 3 and 4. $\square$

Observations. We can extend the above theorems in several ways. First, we can drop the assumption that $Z = Y/\mathcal{G}$ be compact, but then we need to consider bounded, elliptic elements $A \in \text{Hom}(E, E) + \psi_{\text{inv}}^0(Y; E)$ (or, if $\mathcal{G}$ is a vector bundle,

then $A \in \text{Hom}(E, E) + \psi_{\text{inv}}^0(Y; E)$. Also, in the last two theorems, we can allow operators acting between sections of different vector bundles. This will require to slightly modify the proof of Theorem 4, either by using a smooth version of bivariant K -theory [33], or by using the usual bivariant K -theory after we have taken the norm closures of the various ideals $\mathfrak{I}$ decorated with various indices. We can also further integrate along the fibers of $\mathfrak{g}^* \rightarrow B$ to obtain a cohomological formula with values in $H_c^*(B; \mathcal{O})$, the cohomology with local coefficients in the orientation sheaf of $\mathfrak{g}^* \rightarrow B$.

5. INDEX THEORY ON A SIMPLEX

In this section, we discuss an application of our main index theorem to the question of formulating (*i.e.*, the existence of) Fredholm boundary conditions for b -pseudodifferential operators on the simplex

$$\Delta_n = \{(x_0, x_1, \dots, x_n), x_i \geq 0, \sum x_i = 1\}$$

and on manifolds with corners of the form $\Delta_n \times Y$, where Y is a smooth, compact manifold without corners. We obtain complete results for $\dim Y > 0$. The answer to this existence question is “local,” that is, it can be given in terms of the principal symbol. The problem of computing the index of the resulting Fredholm operators, if any, is a “non-local” problem. Some results of this section were obtained in a conversation with R. Melrose, whose insights I am happy to acknowledge.

In addition to the results of the previous sections, we shall also use computations from [28]. All the definitions not included in this section can be found in that paper.

We formalize the above question as the following natural problem:

Problem ($\mathcal{FT}$). *Suppose an elliptic operator $T \in \Psi_b^m(\Delta_n \times Y; E)$ is given. For what T can we find a perturbation $T + R$, with $R \in \Psi_b^{-\infty}(\Delta_n)$, such that*

$$T + R : H^s(\Delta_n \times Y; E) \rightarrow H^{s-m}(\Delta_n \times Y; E)$$

is Fredholm?

We can reduce the general case to the case $m = 0$ by replacing T with $D^{-m}T$, where D is an elliptic, strictly positive operator in $\Psi_b^1(\Delta_n \times Y; E)$. (The resulting operator $D^{-m}T$ is then in the norm completion of $\Psi_b^0(\Delta_n \times Y; \text{End}(E))$, but this makes no difference to us.) For $m = 0$, we then obtain an operator acting on the Hilbert space $L^2(\Delta_n)$.

The answer to the above problem, Problem ($\mathcal{FT}$), is independent on the metric chosen on $\Delta_n \times Y$, although the choice of R may depend on the metric.

Let $\overline{\Psi}_b^0(\Delta_n \times Y)$ be the norm closure of $\Psi_b^0(\Delta_n \times Y)$. By the results of [28], the algebra $\overline{\Psi}_b^0(\Delta_n \times Y)$ contains the ideal of compact operators $\mathcal{K}$ on $L^2(\Delta_n \times Y)$. (See also [18].) Denote by Q be the algebra of “joint symbols” of $\Psi_b^0(\Delta_n \times Y)$:

$$Q := \overline{\Psi}_b^m(\Delta_n \times Y) / \mathcal{K},$$

The principal symbol then extends to a surjective morphism $\sigma_0 : Q \rightarrow C(bS^* \Delta_n \times Y)$ with kernel J .

A first, tautological observation is that Problem ($\mathcal{F}(T)$) has a solution if, and only if, the invertible element $\sigma_0(T) \in C(bS^* \Delta_n \times Y; \text{End}(E))$ has a lifting to an invertible element in Q (the lifting is $T + R$). Since there is a necessary condition for this to happen in terms of K -theory, and the K -groups involved do not depend on E , up to isomorphism, we obtain the following necessary condition for our problem

to have a solution: Problem $(\mathcal{F}(T))$ has a solution only if the class $[\sigma_0(T)] \in K_1(C({}^bS^*\Delta_n \times Y))$ is in the image of the morphism $K_1(Q) \rightarrow K_1(C({}^bS^*\Delta_n \times Y))$.

Now the standard six term K -theory exact sequence associated to the short exact sequence of C_r^* -algebras

$$0 \rightarrow J \rightarrow Q \rightarrow C({}^bS^*\Delta_n \times Y) \rightarrow 0$$

tells us that the class $[\sigma_0(T)] \in K_1(C({}^bS^*\Delta_n \times Y))$ is in the image of the morphism $K_1(Q) \rightarrow K_1(C({}^bS^*\Delta_n \times Y))$ if, and only if, it is in the kernel of the boundary map $\partial : K_1(C({}^bS^*(\Delta_n \times Y))) \rightarrow K_0(J)$ associated to the above exact sequence of algebras.

Lemma 4. *If Problem $(\mathcal{F}_T)$ has a solution, then $\partial[\sigma_m(T)] = 0 \in K_0(J)$. This condition is also sufficient if $\dim Y > 0$.*

Proof. The first part was already proved. The second part follows using the same reasoning as in the proof of from Theorem 1. $\square$

Denote by $I = \overline{\Psi}_b^{-1}(\Delta_n \times Y)$ the norm closure of $\Psi_b^{-1}(\Delta_n \times Y)$ (the closure is in the norm topology of bounded operators on $L^2(\Delta_n \times Y)$). Then $J = I/\mathcal{K}$. The results of [28] give a spectral sequence converging to the K -theory groups of I (and, with some obvious changes, a spectral sequence converging to the K -theory of J). The E^1 -term of this spectral sequence is independent of Y and is (dual to) the combinatorial simplicial complex of Δ_n , more precisely, it is a direct sum of complexes isomorphic to

$$0 \longrightarrow \mathbb{Z}^n \longrightarrow \mathbb{Z}^{\frac{n(n-1)}{2}} \longrightarrow \dots \mathbb{Z}^n \longrightarrow \mathbb{Z} \longrightarrow 0.$$

The direct sum comes from the fact that K -theory is indexed by $\mathbb{Z}/2\mathbb{Z}$ and not by $\mathbb{Z}$. The resulting spectral sequence, which is indexed by $\mathbb{Z} \times \mathbb{Z}$, will be periodic of period 2 in each variable.

It follows that the spectral sequence converging to $K_n(I)$ degenerates at E^2 , which shows that $K_n(I) \simeq \mathbb{Z}$ and $K_{n-1}(I) = 0$ (the same n as in Δ_n). To obtain the K -theory of J one proceeds similarly, the only difference being that one drops the last copy of $\mathbb{Z}$ in the above complex. Finally, this gives $K_0(J) \simeq K_0(I)$ and $K_1(J) \simeq K_1(I) \oplus \mathbb{Z}$.

Denote by

$$\text{In}_k : \Psi_b^\infty(\Delta_n \times Y) \rightarrow \Psi_{\text{inv}}^\infty(Y),$$

$k = 0, \dots, n$, the indicial maps corresponding to the $(n+1)$ -corners of $\Delta_n \times Y$. If B is reduced to a point and $\mathcal{G} = \mathbb{R}^n$, we shall denote $\deg_{\mathcal{G}} = \deg_n$. We are now ready to formulate and prove the main result of this section.

Theorem 6. *Let $T \in \Psi_b^m(\Delta_n \times Y)$, $m > 0$, be an elliptic operator. Assume $\dim Y > 0$. Then, for n odd, Problem $(\mathcal{F}(T))$ always has a solution. For n even, $\deg_n(\text{In}_k(T))$ is independent of k and Problem $(\mathcal{F}(T))$ has a solution if, and only if, $\deg_n(\text{In}_k(T)) = 0$.*

Proof. Using the results of [28], we first observe that, in order for $T + R$ to be Fredholm, it is necessary that $\text{In}_k(T + R)$ be invertible for all k , so

$$\deg_n(\text{In}_k(T)) = \deg_n(\text{In}_k(T + R)) = 0, \quad \text{for all } k.$$

The vanishing of all $\deg_n(\text{In}_k(T))$ is hence necessary for Problem $(\mathcal{F}(T))$ to have a solution. This condition is automatically satisfied for n odd.

To prove that the vanishing of any of $\deg_n(\text{In}_k(T))$ is also enough for Problem $(\mathcal{F}(T))$ to have a solution, we shall show that all degrees $\deg_n(\text{In}_k(T))$ are equal, $k = 1, \dots, n$, and that $\partial[\sigma_0(T)] = 0$ if, and only if, $\deg_n(\text{In}_k(T)) = 0$. The result will follow then from Lemma 4.

As discussed above, it is enough to consider the case $m = 0$. Now

$$K_0(J) = E_{0n}^2 = \ker(\mathbb{Z}^n \rightarrow \mathbb{Z}^{n(n-1)/2}),$$

so $K_0(J) = \{(p, p, \dots, p)\} \subset \mathbb{Z}^n$. Because

$$\partial[\sigma_0(T)] = (\deg_n(\text{In}_0(T)), \deg_n(\text{In}_1(T)), \dots, \deg_n(\text{In}_n(T))) \in E_{0n}^0,$$

we obtain that the degrees $\deg_n(\text{In}_k(T))$ are all equal and that the vanishing of $\partial[\sigma_0(T)]$ is equivalent to the vanishing of any of the degrees $\deg_n(\text{In}_k(T))$. As observed above, an application of Lemma 4 is now enough to complete the proof. $\square$

REFERENCES

- [1] S. Agmon, *On the eigenfunctions and on the eigenvalues of general elliptic boundary value problems*, Comm. Pure Appl. Math. **15** (1962), 627–662.
- [2] B. Ammann, R. Lauter, and V. Nistor, *On the analysis of differential operators on manifolds with a uniform structure at infinity*, work in progress.
- [3] M. F. Atiyah, *K-Theory*, W. A. Benjamin, New York, 1967.
- [4] M. F. Atiyah, *Elliptic operators, discrete subgroups, and von Neumann algebras*, *Astérisque*, **32/33** (1976), 43–72.
- [5] M. F. Atiyah and R. Bott, *The index theorem for manifolds with boundary*, Appendix I in seminar on the Atiyah-Singer index theorem, Annals. of Math. Stud. **57**, Princeton Univ. Press (1965), 337–351.
- [6] M.F. Atiyah, V.K. Patodi, and I.M. Singer, *Spectral asymmetry and Riemannian geometry*, I, Math. Proc. Camb. Phil. Soc. **77** (1975), 43–69.
- [7] M. F. Atiyah and I. M. Singer, *The index of elliptic operators: I*, Ann. of Math. **87** (1968), 484–530.
- [8] M.F. Atiyah and I.M. Singer, *The index of elliptic operators, IV*, Ann. of Math. **93** (1971), 119–138.
- [9] J.-L. Bismut, *Local index theory, eta invariants and holomorphic torsion: a survey*, Surveys in differential geometry, Vol. III (Cambridge, MA, 1996), 1–76, Int. Press, Boston, MA, 1998.
- [10] J.-L. Bismut and J. Cheeger, *Eta invariants and their adiabatic limits*, J. Amer. Math. Soc. **2** (1989), 33–77.
- [11] J.-L. Bismut and D. Freed, *The analysis of elliptic families II. Dirac operators, eta invariants, and the holonomy theorem*, Comm. Math. Phys. **107** (1986), 103–163.
- [12] B. Blackadar, *K-theory for operator algebras*, Cambridge, Cambridge University Press, MSRI Series Publications, no. 5.
- [13] A. Connes, *An analogue of the Thom isomorphism for crossed products of a C^* -algebra by an action of $\mathbb{R}$* , Adv. Math. **39** (1981), 31–55.
- [14] A. Connes, *Noncommutative Geometry*, Academic Press, New York–London, 1994.
- [15] J. Dixmier, *Les C^* -algèbres et leurs représentations*, Gauthier Villars, Paris, 1964.
- [16] G. Grubb and R. Seeley, *Weakly parametric pseudodifferential operators and Atiyah-Patodi-Singer boundary problems*, Inv. Math. **121** (1995), 481–529.
- [17] L. Hörmander, *The analysis of linear partial differential operators*, vol. 3, Springer-Verlag, Berlin, Heidelberg, New York, Tokyo, 1985.
- [18] R. Lauter, B. Monthubert, and V. Nistor, *Pseudodifferential analysis on continuous family groupoids*, Documenta Math. **5** (2000), 625–655 (electronic).
- [19] R. Lauter and S. Moroianu, *Fredholm theory for degenerate pseudodifferential operators on manifolds with fibred boundaries*, to appear in Comm. Partial Diff. Eq.
- [20] R. Lauter and V. Nistor, *Pseudodifferential analysis on groupoids and singular spaces*, Mainz preprint 1999.
- [21] R. Lauter and V. Nistor, *Analysis on open manifolds: A groupoid approach*, Preprint 2000, to appear.

- [22] M. Lesch and M. J. Pflaum, *Traces on algebras of parameter dependent pseudodifferential operators and the eta-invariant*, to appear.
- [23] R. R. Mazzeo, *Elliptic theory of differential edge operators. I*, Comm. Partial Differential Equations, **16**(1991), 1615–1664.
- [24] R. Mazzeo and R.B. Melrose, *Pseudodifferential operators on manifolds with fibered boundaries*. Mikio Sato: a great Japanese mathematician of the twentieth century, Asian J. Math. **2** (1998), 833–866.
- [25] R.B. Melrose, *Pseudodifferential operators, corners, and singular limits*, Proceedings ICM 1990, 217–234, 1991.
- [26] R. B. Melrose, *The Atiyah-Patodi-Singer index theorem*, A. K. Peters, Wellesley, Mass, 1993.
- [27] R. B. Melrose, *Geometric scattering theory*, Stanford Lectures, Cambridge University Press, Cambridge, 1995.
- [28] R. B. Melrose and V. Nistor, *K-theory of C^* -algebras of b-pseudodifferential operators*, GAFA (1998), 88–122.
- [29] R. B. Melrose and P. Piazza, *Families of Dirac operators, boundaries and the b-calculus*, J. Differential Geom. **46** (1997), 99–180.
- [30] S. Moroianu, *Adiabatic limits of eta invariants*, preprint 2001.
- [31] W. Müller, *The trace class conjecture in the theory of automorphic forms*, Ann. of Math. **130** (1989), 605–674.
- [32] W. Müller, *On the analytic continuation of rank one Eisenstein series*, Geom. Funct. Anal. **6** (1996), 572–586.
- [33] V. Nistor, *A bivariate Chern character for p-summable quasi-homomorphisms*, K-theory **5** (1991), 193–211.
- [34] V. Nistor, *A bivariate Chern-Connes character*, Annals. of Math. **138** (1993), 555–590.
- [35] V. Nistor, *On the kernel of the equivariant Dirac operator*, Ann. Global Anal. Geom. **17** (1999), 595–613.
- [36] V. Nistor and E. Troitsky, *A gauge-equivariant index for families of compact groups (tentative title)*, work in progress.
- [37] V. Nistor and A. Weinstein, and Ping Xu, *Pseudodifferential operators on groupoids*, Pacific J. Math. **189** (1999), 117–152.
- [38] S. Sakai, *C^* -algebras and W^* -algebras*, reprint of the 1971 edition. *Classics in Mathematics*, Springer-Verlag, Berlin, 1998, xii+256.
- [39] S. Sethi and M. Stern, *D-brane bound state redux*, Comm. Math. Phys. **194** (1998), 675–705.
- [40] I.M. Singer, *Some remarks on operator theory and index theory*, K-theory and Operator Algebras, Athens, Georgia 1975, (Morrel and Singer, eds.), Lecture Notes in Mathematics, **575**, (1977), 128–138.
- [41] M. A. Shubin, *Pseudodifferential operators and spectral theory*, Springer Verlag, Berlin-Heidelberg-New York, 1987.
- [42] M. E. Taylor, *Pseudodifferential operators*, Princeton Mathematical Series, 34, Princeton University Press, Princeton, N.J., 1981 xi+452.
- [43] M. E. Taylor, *Partial differential equations*, Applied Mathematical Sciences, vol. I-III, Springer-Verlag, New York, (1995-1997).
- [44] E. Troitsky, *Action of compact groups, C^* -index theorem, and families*, math.OA/9904002.
- [45] E. Witten, *Global gravitational anomalies*, Comm. Math. Phys. **100** (1985), 197–229.

CURRENT ADDRESS: DEPARTMENT OF MATHEMATICS, PENNSYLVANIA STATE UNIVERSITY
E-mail address: `nistor@math.psu.edu`